
Trotter-Suzuki-MPI Python Documentation

Release 1.6.1

Peter Wittek, Luca Calderaro

Dec 21, 2016

CONTENTS

1	Introduction	1
1.1	Copyright and License	1
1.2	Acknowledgement	1
1.3	Citations	2
2	Download and Installation	3
2.1	Dependencies	3
3	Quick Start Guide	5
3.1	Simulation Set up	5
3.2	Analysis	6
4	Examples	9
4.1	Expectation values of the Hamiltonian and kinetic operators	9
4.2	Imaginary time evolution to approximate the ground-state energy	10
4.3	Imprinting of a vortex in a Bose-Einstein Condensate	10
4.4	Dark Soliton Generation in Bose-Einstein Condensate using Phase Imprinting	11
5	Function Reference	13
5.1	Lattice1D Class	13
5.2	Lattice2D Class	14
5.3	State Classes	15
5.4	Potential Classes	27
5.5	Hamiltonian Classes	29
5.6	Solver Class	31
5.7	Tools	33
	Index	35

INTRODUCTION

The module is a massively parallel implementation of the Trotter-Suzuki approximation to simulate the evolution of quantum systems classically. It relies on interfacing with C++ code with OpenMP for multicore execution, and it can be accelerated by CUDA.

Key features of the Python interface:

- Simulation of 1D and 2D quantum systems.
- Fast execution by parallelization: OpenMP and CUDA are supported.
- Many-body simulations with non-interacting particles.
- Solving the Gross-Pitaevskii equation (e.g., [dark solitons](#), [vortex dynamics in Bose-Einstein Condensates](#)).
- Imaginary time evolution to approximate the ground state.
- Stationary and time-dependent external potential.
- NumPy arrays are supported for efficient data exchange.
- Multi-platform: Linux, OS X, and Windows are supported.

Copyright and License

Trotter-Suzuki-MPI is free software; you can redistribute it and/or modify it under the terms of the [GNU General Public License](#) as published by the Free Software Foundation; either version 3 of the License, or (at your option) any later version.

Trotter-Suzuki-MPI is distributed in the hope that it will be useful, but WITHOUT ANY WARRANTY; without even the implied warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE. See the [GNU General Public License](#) for more details.

Acknowledgement

The [original high-performance kernels](#) were developed by Carlos Bederián. The distributed extension was carried out while Peter Wittek was visiting the [Department of Computer Applications in Science & Engineering](#) at the [Barcelona Supercomputing Center](#), funded by the “Access to BSC Facilities” project of the [HPC-Europe2](#) programme (contract no. 228398). Generalizing the capabilities of kernels was carried out by Luca Calderaro while visiting the [Quantum Information Theory Group](#) at [ICFO-The Institute of Photonic Sciences](#), sponsored by the [Erasmus+](#) programme. Further computational resources were granted by the [Spanish Supercomputing Network](#) (FY-2015-2-0023 and FI-2016-3-0042) and the [Swedish National Infrastructure for Computing](#) (SNIC 2015/1-162 and 2016/1-320), and a hardware grant by [Nvidia](#). [Pietro Massignan](#) has contributed to the project with extensive testing and suggestions of new features.

Citations

1. Bederián, C. & Dente, A. (2011). Boosting quantum evolutions using Trotter-Suzuki algorithms on GPUs. *Proceedings of HPCLatAm-11, 4th High-Performance Computing Symposium*. [PDF](#)
2. Wittek, P. and Cucchietti, F.M. (2013). [A Second-Order Distributed Trotter-Suzuki Solver with a Hybrid CPU-GPU Kernel](#). *Computer Physics Communications*, 184, pp. 1165-1171. [PDF](#)
3. Wittek, P. and Calderaro, L. (2015). [Extended computational kernels in a massively parallel implementation of the Trotter-Suzuki approximation](#). *Computer Physics Communications*, 197, pp. 339-340. [PDF](#)

DOWNLOAD AND INSTALLATION

The entire package for is available from the [Python Package Index](#), containing the source code and examples. The documentation is hosted on [Read the Docs](#).

The latest development version is available on [GitHub](#). Further examples are available in Jupyter [notebooks](#).

Dependencies

The module requires [Numpy](#). The code is compatible with both Python 2 and 3.

Installation

The code is available on PyPI, hence it can be installed by

```
$ sudo pip install trottersuzuki
```

If you want the latest git version, first clone the repository and generate the Python version. Note that it requires autotools and swig.

```
$ git clone https://github.com/trotter-suzuki-mpi/trotter-suzuki-mpi
$ cd trotter-suzuki-mpi
$ ./autogen.sh
$ ./configure --without-mpi --without-cuda
$ make python
```

Then follow the standard procedure for installing Python modules from the *src/Python* folder:

```
:: $ sudo python setup.py install
```

To install with CUDA support, set the *CUDAHOME* environment variable before calling the Python install.

Build on Mac OS X

Before installing using pip, gcc should be installed first. As of OS X 10.9, gcc is just symlink to clang. To build trottersuzuki and this extension correctly, it is recommended to install gcc using something like:

```
$ brew install gcc48
```

and set environment using:

```
export CC=/usr/local/bin/gcc
export CXX=/usr/local/bin/g++
export CPP=/usr/local/bin/cpp
export LD=/usr/local/bin/gcc
alias c++=/usr/local/bin/c++
alias g++=/usr/local/bin/g++
alias gcc=/usr/local/bin/gcc
alias cpp=/usr/local/bin/cpp
alias ld=/usr/local/bin/gcc
alias cc=/usr/local/bin/gcc
```

Then you can issue

```
$ sudo pip install trottersuzuki
```


QUICK START GUIDE

Simulation Set up

Start by importing the module:

```
import trottersuzuki as ts
```

To set up the simulation of a quantum system, we need only a few lines of code. First of all, we create the lattice over which the physical system is defined. All information about the discretized space is collected in a single object. Say we want a squared lattice of 300x300 nodes, with a physical area of 20x20, then we have to specify these in the constructor of the `Lattice` class:

```
grid = ts.Lattice2D(300, 20.)
```

The object `grid` defines the geometry of the system and it will be used throughout the simulations. Note that the origin of the lattice is at its centre and the lattice points are scaled to the physical locations. The actual transformation is exposed to the user via the function `center_coordinates`.

The physics of the problem is described by the Hamiltonian. A single object is going to store all the information regarding the Hamiltonian. The module is able to deal with two physical models: Gross-Pitaevskii equation of a single or two-component wave function, namely (in units $\hbar = 1$):

$$i \frac{\partial}{\partial t} \psi(t) = H \psi(t)$$

being

$$H = \frac{1}{2m} (P_x^2 + P_y^2) + V(x, y) + g |\psi(x, y)|^2 + \omega L_z$$

and $\psi(t) = \psi_t(x, y)$ for the single component wave function, or

$$H = \begin{bmatrix} H_1 & \frac{\Omega}{2} \\ \frac{\Omega}{2} & H_2 \end{bmatrix}$$

where

$$H_1 = \frac{1}{2m_1} (P_x^2 + P_y^2) + V_1(x, y) + g_1 |\psi(x, y)_1|^2 + g_{12} |\psi(x, y)_2|^2 + \omega L_z$$
$$H_2 = \frac{1}{2m_2} (P_x^2 + P_y^2) + V_2(x, y) + g_2 |\psi(x, y)_2|^2 + g_{12} |\psi(x, y)_1|^2 + \omega L_z$$

and $\psi(t) = \begin{bmatrix} \psi_1(t) \\ \psi_2(t) \end{bmatrix}$, for the two component wave function.

First we define the object for the external potential $V(x, y)$. A general external potential function can be defined by a Python function, for instance, the harmonic potential can be defined as follows:

```
def harmonic_potential(x,y):  
    return 0.5 * (x**2 + y**2)
```

Now we create the external potential object using the `Potential` class and then we initialize it with the function above:

```
potential = ts.Potential(grid) # Create the potential object  
potential.init_potential(harmonic_potential) # Initialize it using a python function
```

Note that the module provides a quick way to define the harmonic potential, as it is frequently used:

```
omegax = omegay = 1.  
harmonicpotential = ts.HarmonicPotential(grid, omegax, omegay)
```

We are ready to create the Hamiltonian object. For the sake of simplicity, let us create the Hamiltonian of the harmonic oscillator:

```
particle_mass = 1. # Mass of the particle  
hamiltonian = ts.Hamiltonian(grid, potential, particle_mass) # Create the_  
↪Hamiltonian object
```

The quantum state is created by the `State` class; it resembles the way the potential is defined. Here we create the ground state of the harmonic oscillator:

```
import numpy as np # Import the module numpy for the exponential and sqrt functions  
  
def state_wave_function(x,y): # Wave function  
    return np.exp(-0.5*(x**2 + y**2)) / np.sqrt(np.pi)  
  
state = ts.State(grid) # Create the quantum state  
state.init_state(state_wave_function) # Initialize the state
```

The module provides several predefined quantum states as well. In this case, we could have used the `GaussianState` class:

```
omega = 1.  
gaussianstate = ts.GaussianState(grid, omega) # Create a quantum state whose wave_  
↪function is Gaussian-like
```

We are left with the creation of the last object: the `Solver` class gathers all the objects we defined so far and it is used to perform the evolution and analyze the expectation values:

```
delta_t = 1e-3 # Physical time of a single iteration  
solver = ts.Solver(grid, state, hamiltonian, delta_t) # Creating the solver object
```

Finally we can perform both real-time and imaginary-time evolution using the method `evolve`:

```
iterations = 100 # Number of iterations to be performed  
solver.evolve(iterations, True) # Perform imaginary-time evolution  
solver.evolve(iterations) # Perform real-time evolution
```

Analysis

The classes we have seen so far implement several members useful to analyze the system (see the function reference section for a complete list).

Expectation values

The solver class provides members for the energy calculations. For instance, the total energy can be calculated using the `get_total_energy` member. We expect it to be 1 ($\hbar = 1$), and indeed we get the right result up to a small error which depends on the lattice approximation:

```
tot_energy = solver.get_total_energy()
print(tot_energy)
```

```
1.00146456951
```

The expected values of the X , Y , P_x , P_y operators are calculated using the members in the `State` class

```
mean_x = state.get_mean_x()  # Get the expected value of X operator
print(mean_x)
```

```
1.39431975344e-14
```

Norm of the state

The squared norm of the state can be calculated by means of both `State` and `Solver` classes

```
snorm = state.get_squared_norm()
print(snorm)
```

```
1.0
```

Particle density and Phase

Very often one is interested in the phase and particle density of the state. Two members of `State` class provide these features

```
density = state.get_particle_density()  # Return a numpy matrix of the particle_
↪density
phase = state.get_phase()  # Return a numpy matrix of the phase
```

Imprinting

The member `imprint`, in the `State` class, applies the following transformation to the state:

$$\psi(x, y) \rightarrow \psi'(x, y) = f(x, y)\psi(x, y)$$

being $f(x, y)$ a general complex-valued function. This comes in handy when we want to imprint, for instance, vortices or solitons:

```
def vortex(x, y):  # Function defining a vortex
    z = x + 1j*y
    angle = np.angle(z)
    return np.exp(1j * angle)

state.imprint(vortex)  # Imprint the vortex on the state
```

File Input and Output

`write_to_files` and `loadtxt` members, in `State` class, provide a simple way to handle file I/O. The former writes the wave function arranged as a complex matrix, in a plain text; the latter loads the wave function from a file to the state object. The following code provides an example:

```
state.write_to_file("file_name") # Write the wave function to a file
state2 = ts.State(grid) # Create a new state
state2.loadtxt("file_name") # Load the wave function from the file
```

For a complete list of methods see the function reference.

EXAMPLES

Expectation values of the Hamiltonian and kinetic operators

The following code block gives a simple example of initializing a state and calculating the expectation values of the Hamiltonian and kinetic operators and the norm of the state after the evolution.

```
import numpy as np
from trottersuzuki import *

grid = Lattice2D(256, 15) # create a 2D lattice

potential = HarmonicPotential(grid, 1, 1) # define an symmetric harmonic potential_
↪with unit frequency
particle_mass = 1.
hamiltonian = Hamiltonian(grid, potential, particle_mass) # define the_
↪Hamiltonian:

frequency = 1
state = GaussianState(grid, frequency) # define gaussian wave function state: we_
↪choose the ground state of the Hamiltonian

time_of_single_iteration = 1.e-4
solver = Solver(grid, state, hamiltonian, time_of_single_iteration) # define the_
↪solver

# get some expected values from the initial state
print("norm: ", solver.get_squared_norm())
print("Total energy: ", solver.get_total_energy())
print("Kinetic energy: ", solver.get_kinetic_energy())

number_of_iterations = 1000
solver.evolve(number_of_iterations) # evolve the state of 1000 iterations

# get some expected values from the evolved state
print("norm: ", solver.get_squared_norm())
print("Total energy: ", solver.get_total_energy())
print("Kinetic energy: ", solver.get_kinetic_energy())
```

Imaginary time evolution to approximate the ground-state energy

```
import numpy as np
from trottersuzuki import *

grid = Lattice2D(256, 15) # create a 2D lattice

potential = HarmonicPotential(grid, 1, 1) # define an symmetric harmonic potential_
↳with unit frequency
particle_mass = 1.
hamiltonian = Hamiltonian(grid, potential, particle_mass) # define the Hamiltonian:

frequency = 3
state = GaussianState(grid, frequency) # define gaussian wave function state: we_
↳choose the ground state of the Hamiltonian

time_of_single_iteration = 1.e-4
solver = Solver(grid, state, hamiltonian, time_of_single_iteration) # define the_
↳solver

# get some expected values from the initial state
print("norm: ", solver.get_squared_norm())
print("Total energy: ", solver.get_total_energy())
print("Kinetic energy: ", solver.get_kinetic_energy())

number_of_iterations = 40000
imaginary_evolution = True
solver.evolve(number_of_iterations, imaginary_evolution) # evolve the state of_
↳40000 iterations

# get some expected values from the evolved state
print("norm: ", solver.get_squared_norm())
print("Total energy: ", solver.get_total_energy())
print("Kinetic energy: ", solver.get_kinetic_energy())
```

Imprinting of a vortex in a Bose-Einstein Condensate

```
import numpy as np
import trottersuzuki as ts

grid = ts.Lattice2D(256, 15) # create a 2D lattice

potential = HarmonicPotential(grid, 1, 1) # define an symmetric harmonic potential_
↳with unit frequency
particle_mass = 1.
coupling_intra_particle_interaction = 100.
hamiltonian = Hamiltonian(grid, potential, particle_mass, coupling_intra_particle_
↳interaction) # define the Hamiltonian:

frequency = 1
state = GaussianState(grid, frequency) # define gaussian wave function state: we_
↳choose the ground state of the Hamiltonian
def vortex(x, y): # vortex to be imprinted
    z = x + 1j*y
    angle = np.angle(z)
```

```

    return np.exp(1j * angle)

state.imprint(vortex) # imprint the vortex on the condensate

time_of_single_iteration = 1.e-4
solver = Solver(grid, state, hamiltonian, time_of_single_iteration) # define the_
↪solver

```

Dark Soliton Generation in Bose-Einstein Condensate using Phase Imprinting

This example simulates the evolution of a dark soliton in a Bose-Einstein Condensate. For a more detailed description, refer to [this notebook](#).

```

from __future__ import print_function
import numpy as np
import trottersuzuki as ts
from matplotlib import pyplot as plt

grid = ts.Lattice2D(300, 50.) # # create a 2D lattice

potential = ts.HarmonicPotential(grid, 1., 1./np.sqrt(2.)) # create an harmonic_
↪potential
coupling = 1.2097e3
hamiltonian = ts.Hamiltonian(grid, potential, 1., coupling) # create the Hamiltonian

state = ts.GaussianState(grid, 0.05) # create the initial state
solver = ts.Solver(grid, state, hamiltonian, 1.e-4) # initialize the solver
solver.evolve(10000, True) # evolve the state towards the ground state

density = state.get_particle_density()
plt.pcolor(density) # plot the particle denisity
plt.show()



def dark_soliton(x,y): # define phase imprinting that will create the dark soliton
    a = 1.98128
    theta = 1.5*np.pi
    return np.exp(1j* (theta * 0.5 * (1. + np.tanh(-a * x))))

state.imprint(dark_soliton) # phase imprinting
solver.evolve(1000) # perform a real time evolution

density = state.get_particle_density()
plt.pcolor(density) # plot the particle denisity
plt.show()

```

The results are the following plots:



FUNCTION REFERENCE

Lattice1D Class

class trottersuzuki.**Lattice1D**

This class defines the lattice structure over which the state and potential matrices are defined.

Constructors

Lattice2D (*dim_x*, *length_x*, *periodic_x_axis=False*)

Construct the Lattice2D.

Parameters

- **dim_x** [integer] Linear dimension of the squared lattice in the x direction.
- **length_x** [float] Physical length of the lattice's side in the x direction.
- **periodic_x_axis** [bool, optional (default: False)] Boundary condition along the x axis (false=closed, true=periodic).

Returns

- **Lattice1D** [Lattice1D object] Define the geometry of the simulation.

Notes

The lattice created is squared.

Example

```
>>> import trottersuzuki as ts # import the module
>>> # Generate a 200-point Lattice1D with physical dimensions of 30
>>> # and closed boundary conditions.
>>> grid = ts.Lattice1D(200, 30.)
```

Members

get_x_axis()

Get the x-axis of the lattice.

Returns

- **x_axis** [numpy array] X-axis of the lattice

Attributes

length_x

Physical length of the lattice along the X-axis.

dim_x

Number of dots of the lattice along the X-axis.

delta_x

Resolution of the lattice along the X-axis: ratio between *length_x* and *dim_x*.

Lattice2D Class

class trottersuzuki.**Lattice2D**

This class defines the lattice structure over which the state and potential matrices are defined.

Constructors

Lattice2D(*dim_x*, *length_x*, *dim_y=None*, *length_y=None*, *periodic_x_axis=False*, *periodic_y_axis=False*)
Construct the Lattice2D.

Parameters

- dim_x** [integer] Linear dimension of the squared lattice in the x direction.
- length_x** [float] Physical length of the lattice's side in the x direction.
- dim_y** [integer,optional (default: equal to dim_x)] Linear dimension of the squared lattice in the y direction.
- length_y** [float,optional (default: equal to length_x)] Physical length of the lattice's side in the y direction.
- periodic_x_axis** [bool,optional (default: False)] Boundary condition along the x axis (false=closed, true=periodic).
- periodic_y_axis** [bool,optional (default: False)] Boundary condition along the y axis (false=closed, true=periodic).

Returns

- Lattice2D** [Lattice2D object] Define the geometry of the simulation.

Notes

The lattice created is squared.

Example

```
>>> import trottersuzuki as ts # import the module
>>> # Generate a 200x200 Lattice2D with physical dimensions of 30x30
>>> # and closed boundary conditions.
>>> grid = ts.Lattice2D(200, 30.)
```

Members

get_x_axis()

Get the x-axis of the lattice.

Returns

- x_axis** [numpy array] X-axis of the lattice

get_y_axis()

Get the y-axis of the lattice.

Returns

- y_axis** [numpy array] Y-axis of the lattice

Attributes

length_x

Physical length of the lattice along the X-axis.

length_y

Physical length of the lattice along the Y-axis.

dim_x

Number of dots of the lattice along the X-axis.

dim_y

Number of dots of the lattice along the Y-axis.

delta_x

Resolution of the lattice along the X-axis: ratio between *length_x* and *dim_x*.

delta_y

Resolution of the lattice along the y-axis: ratio between *length_y* and *dim_y*.

State Classes

class trottersuzuki.**State**

This class defines the quantum state.

Constructors

State (*grid*)

Create a quantum state.

Parameters

- grid** [Lattice object] Define the geometry of the simulation.

Returns

- state** [State object] Quantum state.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> def wave_function(x,y): # Define a flat wave function
>>>     return 1.
>>> state = ts.State(grid) # Create the system's state
>>> state.ini_state(wave_function) # Initialize the wave function of the
↪state
```

State (*state*)

Copy a quantum state.

Parameters

- state** [State object] Quantum state to be copied

Returns

- state** [State object] Quantum state.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state with a
↳ gaussian wave function
>>> state2 = ts.State(state) # Copy state into state2
```

Members

State.init_state(state_function) :

Initialize the wave function of the state using a function.

Parameters

•**state_function** [python function] Python function defining the wave function of the state ψ .

Notes

The input arguments of the python function must be (x,y).

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> def wave_function(x,y): # Define a flat wave function
>>>     return 1.
>>> state = ts.State(grid) # Create the system's state
>>> state.init_state(wave_function) # Initialize the wave function of the
↳ state
```

imprint (function)

Multiply the wave function of the state by the function provided.

Parameters

•**function** [python function] Function to be printed on the state.

Notes

Useful, for instance, to imprint solitons and vortices on a condensate. Generally, it performs a transformation of the state whose wave function becomes:

$$\psi(x,y)' = f(x,y)\psi(x,y)$$

being $f(x,y)$ the input function and $\psi(x,y)$ the initial wave function.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> def vortex(x,y): # Vortex function
>>>     z = x + 1j*y
>>>     angle = np.angle(z)
>>>     return np.exp(1j * angle)
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.imprint(vortex) # Imprint a vortex on the state
```

get_mean_px()

Return the expected value of the P_x operator.

Returns

•**mean_px** [float] Expected value of the P_x operator.

get_mean_ppxx()

Return the expected value of the P_x^2 operator.

Returns

•*mean_ppxx* [float] Expected value of the P_x^2 operator.

get_mean_py()

Return the expected value of the P_y operator.

Returns

•*mean_py* [float] Expected value of the P_y operator.

get_mean_pypy()

Return the expected value of the P_y^2 operator.

Returns

•*mean_pypy* [float] Expected value of the P_y^2 operator.

get_mean_x()

Return the expected value of the X operator.

Returns

•*mean_x* [float] Expected value of the X operator.

get_mean_xx()

Return the expected value of the X^2 operator.

Returns

•*mean_xx* [float] Expected value of the X^2 operator.

get_mean_y()

Return the expected value of the Y operator.

Returns

•*mean_y* [float] Expected value of the Y operator.

get_mean_yy()

Return the expected value of the Y^2 operator.

Returns

•*mean_yy* [float] Expected value of the Y^2 operator.

get_particle_density()

Return a matrix storing the squared norm of the wave function.

Returns

•*particle_density* [numpy matrix] Particle density of the state $|\psi(x, y)|^2$

get_phase()

Return a matrix of the wave function's phase.

Returns

•*get_phase* [numpy matrix] Matrix of the wave function's phase $\phi(x, y) = \log(\psi(x, y))$

get_squared_norm()

Return the squared norm of the quantum state.

Returns

•**squared_norm** [float] Squared norm of the quantum state.

loadtxt (*file_name*)

Load the wave function from a file.

Parameters

•**file_name** [string] Name of the file to be written.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.write_to_file('wave_function.txt') # Write to a file the wave_
↪function
>>> state2 = ts.State(grid) # Create a quantum state
>>> state2.loadtxt('wave_function.txt') # Load the wave function
```

write_particle_density (*file_name*)

Write to a file the particle density matrix of the wave function.

Parameters

•**file_name** [string] Name of the file.

write_phase (*file_name*)

Write to a file the wave function.

Parameters

•**file_name** [string] Name of the file to be written.

write_to_file (*file_name*)

Write to a file the wave function.

Parameters

•**file_name** [string] Name of the file to be written.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.write_to_file('wave_function.txt') # Write to a file the wave_
↪function
>>> state2 = ts.State(grid) # Create a quantum state
>>> state2.loadtxt('wave_function.txt') # Load the wave function
```

class trottersuzuki.**ExponentialState**

This class defines a quantum state with exponential like wave function.

This class is a child of State class.

Constructors

ExponentialState (*grid, n_x=1, n_y=1, norm=1, phase=0*)

Construct the quantum state with exponential like wave function.

Parameters

•**grid** [Lattice2D object] Defines the geometry of the simulation.

•**n_x** [integer, optional (default: 1)] First quantum number.

- n_y*** [integer, optional (default: 1)] Second quantum number.
- norm*** [float, optional (default: 1)] Squared norm of the quantum state.
- phase*** [float, optional (default: 0)] Relative phase of the wave function.

Returns

- ExponentialState*** [State object.] Quantum state with exponential like wave function. The wave function is give by:n

$$\psi(x, y) = \sqrt{N}/L e^{i2\pi(n_x x + n_y y)/L} e^{i\phi}$$

being N the norm of the state, L the length of the lattice edge, n_x and n_y the quantum numbers and ϕ the relative phase.

Notes

The geometry of the simulation has to have periodic boundary condition to use Exponential state as initial state of a real time evolution. Indeed, the wave function is not null at the edges of the space.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D(300, 30., True, True) # Define the simulation's
↳ geometry
>>> state = ts.ExponentialState(grid, 2, 1) # Create the system's state
```

Member

imprint (*function*)

Multiply the wave function of the state by the function provided.

Parameters

- function*** [python function] Function to be printed on the state.

Notes

Useful, for instance, to imprint solitons and vortices on a condensate. Generally, it performs a transformation of the state whose wave function becomes:

$$\psi(x, y)' = f(x, y)\psi(x, y)$$

being $f(x, y)$ the input function and $\psi(x, y)$ the initial wave function.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> def vortex(x, y): # Vortex function
>>>     z = x + 1j*y
>>>     angle = np.angle(z)
>>>     return np.exp(1j * angle)
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.imprint(vortex) # Imprint a vortex on the state
```

get_mean_px ()

Return the expected value of the P_x operator.

Returns

- mean_px*** [float] Expected value of the P_x operator.

get_mean_ppx()

Return the expected value of the P_x^2 operator.

Returns

• *mean_ppx* [float] Expected value of the P_x^2 operator.

get_mean_py()

Return the expected value of the P_y operator.

Returns

• *mean_py* [float] Expected value of the P_y operator.

get_mean_pypy()

Return the expected value of the P_y^2 operator.

Returns

• *mean_pypy* [float] Expected value of the P_y^2 operator.

get_mean_x()

Return the expected value of the X operator.

Returns

• *mean_x* [float] Expected value of the X operator.

get_mean_xx()

Return the expected value of the X^2 operator.

Returns

• *mean_xx* [float] Expected value of the X^2 operator.

get_mean_y()

Return the expected value of the Y operator.

Returns

• *mean_y* [float] Expected value of the Y operator.

get_mean_yy()

Return the expected value of the Y^2 operator.

Returns

• *mean_yy* [float] Expected value of the Y^2 operator.

get_particle_density()

Return a matrix storing the squared norm of the wave function.

Returns

• *particle_density* [numpy matrix] Particle density of the state $|\psi(x, y)|^2$

get_phase()

Return a matrix of the wave function's phase.

Returns

• *get_phase* [numpy matrix] Matrix of the wave function's phase $\phi(x, y) = \log(\psi(x, y))$

get_squared_norm()

Return the squared norm of the quantum state.

Returns

•**squared_norm** [float] Squared norm of the quantum state.

loadtxt (*file_name*)

Load the wave function from a file.

Parameters

•**file_name** [string] Name of the file to be written.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.write_to_file('wave_function.txt') # Write to a file the wave_
↪function
>>> state2 = ts.State(grid) # Create a quantum state
>>> state2.loadtxt('wave_function.txt') # Load the wave function
```

write_particle_density (*file_name*)

Write to a file the particle density matrix of the wave function.

Parameters

•**file_name** [string] Name of the file.

write_phase (*file_name*)

Write to a file the wave function.

Parameters

•**file_name** [string] Name of the file to be written.

write_to_file (*file_name*)

Write to a file the wave function.

Parameters

•**file_name** [string] Name of the file to be written.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.write_to_file('wave_function.txt') # Write to a file the wave_
↪function
>>> state2 = ts.State(grid) # Create a quantum state
>>> state2.loadtxt('wave_function.txt') # Load the wave function
```

class trottersuzuki.**GaussianState**

This class defines a quantum state with gaussian like wave function.

This class is a child of State class.

Constructors

GaussianState (*grid, omega_x, omega_y=omega_x, mean_x=0, mean_y=0, norm=1, phase=0*)

Construct the quantum state with gaussian like wave function.

Parameters

•**grid** [Lattice2D object] Defines the geometry of the simulation.

•**omega_x** [float] Inverse of the variance along x-axis.

- omega_y** [float, optional (default: omega_x)] Inverse of the variance along y-axis.
- mean_x** [float, optional (default: 0)] X coordinate of the gaussian function's peak.
- mean_y** [float, optional (default: 0)] Y coordinate of the gaussian function's peak.
- norm** [float, optional (default: 1)] Squared norm of the state.
- phase** [float, optional (default: 0)] Relative phase of the wave function.

Returns

- GaussianState** [State object.] Quantum state with gaussian like wave function. The wave function is given by:

$$\psi(x, y) = (N/\pi)^{1/2} (\omega_x \omega_y)^{1/4} e^{-(\omega_x (x - \mu_x)^2 + \omega_y (y - \mu_y)^2)/2} e^{i\phi}$$

being N the norm of the state, ω_x and ω_y the inverse of the variances, μ_x and μ_y the coordinates of the function's peak and ϕ the relative phase.

Notes

The physical dimensions of the Lattice2D have to be enough to ensure that the wave function is almost zero at the edges.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D(300, 30.) # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 2.) # Create the system's state
```

Members

imprint (function)

Multiply the wave function of the state by the function provided.

Parameters

- function** [python function] Function to be printed on the state.

Notes

Useful, for instance, to imprint solitons and vortices on a condensate. Generally, it performs a transformation of the state whose wave function becomes:

$$\psi(x, y)' = f(x, y)\psi(x, y)$$

being $f(x, y)$ the input function and $\psi(x, y)$ the initial wave function.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> def vortex(x, y): # Vortex function
>>>     z = x + 1j*y
>>>     angle = np.angle(z)
>>>     return np.exp(1j * angle)
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.imprint(vortex) # Imprint a vortex on the state
```

get_mean_px()

Return the expected value of the P_x operator.

Returns

•*mean_px* [float] Expected value of the P_x operator.

get_mean_ppx ()

Return the expected value of the P_x^2 operator.

Returns

•*mean_ppx* [float] Expected value of the P_x^2 operator.

get_mean_py ()

Return the expected value of the P_y operator.

Returns

•*mean_py* [float] Expected value of the P_y operator.

get_mean_pypy ()

Return the expected value of the P_y^2 operator.

Returns

•*mean_pypy* [float] Expected value of the P_y^2 operator.

get_mean_x ()

Return the expected value of the X operator.

Returns

•*mean_x* [float] Expected value of the X operator.

get_mean_xx ()

Return the expected value of the X^2 operator.

Returns

•*mean_xx* [float] Expected value of the X^2 operator.

get_mean_y ()

Return the expected value of the Y operator.

Returns

•*mean_y* [float] Expected value of the Y operator.

get_mean_yy ()

Return the expected value of the Y^2 operator.

Returns

•*mean_yy* [float] Expected value of the Y^2 operator.

get_particle_density ()

Return a matrix storing the squared norm of the wave function.

Returns

•*particle_density* [numpy matrix] Particle density of the state $|\psi(x, y)|^2$

get_phase ()

Return a matrix of the wave function's phase.

Returns

•*get_phase* [numpy matrix] Matrix of the wave function's phase $\phi(x, y) = \log(\psi(x, y))$

get_squared_norm()

Return the squared norm of the quantum state.

Returns

•**squared_norm** [float] Squared norm of the quantum state.

loadtxt (*file_name*)

Load the wave function from a file.

Parameters

•**file_name** [string] Name of the file to be written.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.write_to_file('wave_function.txt') # Write to a file the wave_
↪function
>>> state2 = ts.State(grid) # Create a quantum state
>>> state2.loadtxt('wave_function.txt') # Load the wave function
```

write_particle_density (*file_name*)

Write to a file the particle density matrix of the wave function.

Parameters

•**file_name** [string] Name of the file.

write_phase (*file_name*)

Write to a file the wave function.

Parameters

•**file_name** [string] Name of the file to be written.

write_to_file (*file_name*)

Write to a file the wave function.

Parameters

•**file_name** [string] Name of the file to be written.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.write_to_file('wave_function.txt') # Write to a file the wave_
↪function
>>> state2 = ts.State(grid) # Create a quantum state
>>> state2.loadtxt('wave_function.txt') # Load the wave function
```

class trottersuzuki.**SinusoidState**

This class defines a quantum state with sinusoidal like wave function.

This class is a child of State class.

Constructors

SinusoidState (*grid, n_x=1, n_y=1, norm=1, phase=0*)

Construct the quantum state with sinusoidal like wave function.

Parameters

- grid** [Lattice2D object] Define the geometry of the simulation.
- n_x** [integer, optional (default: 1)] First quantum number.
- n_y** [integer, optional (default: 1)] Second quantum number.
- norm** [float, optional (default: 1)] Squared norm of the quantum state.
- phase** [float, optional (default: 1)] Relative phase of the wave function.

Returns

- SinusoidState** [State object.] Quantum state with sinusoidal like wave function. The wave function is given by:

$$\psi(x, y) = 2\sqrt{N}/L \sin(2\pi n_x x/L) \sin(2\pi n_y y/L) e^{i\phi}$$

being N the norm of the state, L the length of the lattice edge, n_x and n_y the quantum numbers and ϕ the relative phase.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D(300, 30., True, True) # Define the simulation's
↳ geometry
>>> state = ts.SinusoidState(grid, 2, 0) # Create the system's state
```

Members**imprint** (function)

Multiply the wave function of the state by the function provided.

Parameters

- function** [python function] Function to be printed on the state.

Notes

Useful, for instance, to imprint solitons and vortices on a condensate. Generally, it performs a transformation of the state whose wave function becomes:

$$\psi(x, y)' = f(x, y)\psi(x, y)$$

being $f(x, y)$ the input function and $\psi(x, y)$ the initial wave function.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> def vortex(x, y): # Vortex function
>>>     z = x + 1j*y
>>>     angle = np.angle(z)
>>>     return np.exp(1j * angle)
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.imprint(vortex) # Imprint a vortex on the state
```

get_mean_px()

Return the expected value of the P_x operator.

Returns

•*mean_px* [float] Expected value of the P_x operator.

get_mean_ppx()

Return the expected value of the P_x^2 operator.

Returns

•*mean_ppx* [float] Expected value of the P_x^2 operator.

get_mean_py()

Return the expected value of the P_y operator.

Returns

•*mean_py* [float] Expected value of the P_y operator.

get_mean_pypy()

Return the expected value of the P_y^2 operator.

Returns

•*mean_pypy* [float] Expected value of the P_y^2 operator.

get_mean_x()

Return the expected value of the X operator.

Returns

•*mean_x* [float] Expected value of the X operator.

get_mean_xx()

Return the expected value of the X^2 operator.

Returns

•*mean_xx* [float] Expected value of the X^2 operator.

get_mean_y()

Return the expected value of the Y operator.

Returns

•*mean_y* [float] Expected value of the Y operator.

get_mean_yy()

Return the expected value of the Y^2 operator.

Returns

•*mean_yy* [float] Expected value of the Y^2 operator.

get_particle_density()

Return a matrix storing the squared norm of the wave function.

Returns

•*particle_density* [numpy matrix] Particle density of the state $|\psi(x, y)|^2$

get_phase()

Return a matrix of the wave function's phase.

Returns

•*get_phase* [numpy matrix] Matrix of the wave function's phase $\phi(x, y) = \log(\psi(x, y))$

get_squared_norm()

Return the squared norm of the quantum state.

Returns

•**squared_norm** [float] Squared norm of the quantum state.

loadtxt (*file_name*)

Load the wave function from a file.

Parameters

•**file_name** [string] Name of the file to be written.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.write_to_file('wave_function.txt') # Write to a file the wave_
↪function
>>> state2 = ts.State(grid) # Create a quantum state
>>> state2.loadtxt('wave_function.txt') # Load the wave function
```

write_particle_density (*file_name*)

Write to a file the particle density matrix of the wave function.

Parameters

•**file_name** [string] Name of the file.

write_phase (*file_name*)

Write to a file the wave function.

Parameters

•**file_name** [string] Name of the file to be written.

write_to_file (*file_name*)

Write to a file the wave function.

Parameters

•**file_name** [string] Name of the file to be written.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> state.write_to_file('wave_function.txt') # Write to a file the wave_
↪function
>>> state2 = ts.State(grid) # Create a quantum state
>>> state2.loadtxt('wave_function.txt') # Load the wave function
```

Potential Classes

class trottersuzuki.**Potential**

This class defines the external potential that is used for Hamiltonian class.

Constructors

Potential (*grid*)

Construct the external potential.

Parameters

•*grid* [Lattice object] Define the geometry of the simulation.

Returns

•*Potential* [Potential object] Create external potential.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> # Define a constant external potential
>>> def external_potential_function(x, y):
>>>     return 1.
>>> potential = ts.Potential(grid) # Create the external potential
>>> potential.init_potential(external_potential_function) # Initialize the_
↪external potential
```

Members**init_potential** (*potential_function*)

Initialize the external potential.

Parameters

•*potential_function* [python function] Define the external potential function.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> # Define a constant external potential
>>> def external_potential_function(x, y):
>>>     return 1.
>>> potential = ts.Potential(grid) # Create the external potential
>>> potential.init_potential(external_potential_function) # Initialize the_
↪external potential
```

get_value (*x, y*)

Get the value at the lattice's coordinate (x,y).

Returns

•*value* [float] Value of the external potential.

class trottersuzuki.**HarmonicPotential**

This class defines the external potential, that is used for Hamiltonian class.

This class is a child of Potential class.

Constructors**HarmonicPotential**(*grid, omegax, omegay, mass=1., mean_x=0., mean_y=0.*)`

Construct the harmonic external potential.

Parameters

•*grid* [Lattice2D object] Define the geometry of the simulation.

•*omegax* [float] Frequency along x-axis.

- omegay** [float] Frequency along y-axis.
- mass** [float,optional (default: 1.)] Mass of the particle.
- mean_x** [float,optional (default: 0.)] Minimum of the potential along x axis.
- mean_y** [float,optional (default: 0.)] Minimum of the potential along y axis.

Returns

- HarmonicPotential** [Potential object] Harmonic external potential.

Notes

External potential function:n

$$V(x, y) = 1/2m(\omega_x^2 x^2 + \omega_y^2 y^2)$$

being m the particle mass, ω_x and ω_y the potential frequencies.

Example

```
>>> import trottersuzuki as ts # Import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> potential = ts.HarmonicPotential(grid, 2., 1.) # Create an harmonic_
↪external potential
```

Members

get_value (x, y)

Get the value at the lattice's coordinate (x, y).

Returns

- value** [float] Value of the external potential.

Hamiltonian Classes

class trottersuzuki.**Hamiltonian**

This class defines the Hamiltonian of a single component system.

Constructors

Hamiltonian ($grid$, $potential=0$, $mass=1.$, $coupling=0.$, $angular_velocity=0.$, $rot_coord_x=0$, $rot_coord_y=0$)

Construct the Hamiltonian of a single component system.

Parameters

- grid** [Lattice object] Define the geometry of the simulation.
- potential** [Potential object] Define the external potential of the Hamiltonian (V).
- mass** [float,optional (default: 1.)] Mass of the particle (m).
- coupling** [float,optional (default: 0.)] Coupling constant of intra-particle interaction (g).
- angular_velocity** [float,optional (default: 0.)] The frame of reference rotates with this angular velocity (ω).
- rot_coord_x** [float,optional (default: 0.)] X coordinate of the center of rotation.
- rot_coord_y** [float,optional (default: 0.)] Y coordinate of the center of rotation.

Returns

- Hamiltonian** [Hamiltonian object] Hamiltonian of the system to be simulated:

$$H(x, y) = \frac{1}{2m}(P_x^2 + P_y^2) + V(x, y) + g|\psi(x, y)|^2 + \omega L_z$$

being m the particle mass, $V(x, y)$ the external potential, g the coupling constant of intra-particle interaction, ω the angular velocity of the frame of reference and L_z the angular momentum operator along the z-axis.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> potential = ts.HarmonicPotential(grid, 1., 1.) # Create an harmonic_
↪external potential
>>> hamiltonian = ts.Hamiltonian(grid, potential) # Create the Hamiltonian_
↪of an harmonic oscillator
```

class trottersuzuki.Hamiltonian2Component

This class defines the Hamiltonian of a two component system.

Constructors

Hamiltonian2Component (*grid*, *potential_1*=0, *potential_2*=0, *mass_1*=1., *mass_2*=1., *coupling_1*=0., *coupling_12*=0., *coupling_2*=0., *omega_r*=0, *omega_i*=0, *angular_velocity*=0., *rot_coord_x*=0, *rot_coord_y*=0)

Construct the Hamiltonian of a two component system.

Parameters

- grid** [Lattice object] Define the geometry of the simulation.
- potential_1** [Potential object] External potential to which the first state is subjected (V_1).
- potential_2** [Potential object] External potential to which the second state is subjected (V_2).
- mass_1** [float, optional (default: 1.)] Mass of the first-component's particles (m_1).
- mass_2** [float, optional (default: 1.)] Mass of the second-component's particles (m_2).
- coupling_1** [float, optional (default: 0.)] Coupling constant of intra-particle interaction for the first component (g_1).
- coupling_12** [float, optional (default: 0.)] Coupling constant of inter-particle interaction between the two components (g_{12}).
- coupling_2** [float, optional (default: 0.)] Coupling constant of intra-particle interaction for the second component (g_2).
- omega_r** [float, optional (default: 0.)] Real part of the Rabi coupling ($\text{Re}(\Omega)$).
- omega_i** [float, optional (default: 0.)] Imaginary part of the Rabi coupling ($\text{Im}(\Omega)$).
- angular_velocity** [float, optional (default: 0.)] The frame of reference rotates with this angular velocity (ω).
- rot_coord_x** [float, optional (default: 0.)] X coordinate of the center of rotation.
- rot_coord_y** [float, optional (default: 0.)] Y coordinate of the center of rotation.

Returns

- Hamiltonian2Component** [Hamiltonian2Component object] Hamiltonian of the two-component system to be simulated.

$$H = \begin{bmatrix} H_1 & \frac{\Omega}{2} \\ \frac{\Omega}{2} & H_2 \end{bmatrix}$$

being

$$H_1 = \frac{1}{2m_1}(P_x^2 + P_y^2) + V_1(x, y) + g_1|\psi_1(x, y)|^2 + g_{12}|\psi_2(x, y)|^2 + \omega L_z$$

$$H_2 = \frac{1}{2m_2}(P_x^2 + P_y^2) + V_2(x, y) + g_2|\psi_2(x, y)|^2 + g_{12}|\psi_1(x, y)|^2 + \omega L_z$$

and, for the i -th component, m_i the particle mass, $V_i(x, y)$ the external potential, g_i the coupling constant of intra-particle interaction; g_{12} the coupling constant of inter-particle interaction ω the angular velocity of the frame of reference, L_z the angular momentum operator along the z -axis and Ω the Rabi coupling.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> potential = ts.HarmonicPotential(grid, 1., 1.) # Create an harmonic_
↪external potential
>>> hamiltonian = ts.Hamiltonian2Component(grid, potential, potential) #_
↪Create the Hamiltonian of an harmonic oscillator for a two-component system
```

Solver Class

class trottersuzuki.**Solver**

This class defines the evolution tasks.

Constructors

Solver(grid, state1, hamiltonian, delta_t, Potential=None, State2=None, Potential2=None, kernel_type="cpu")

Construct the Solver object. Potential is only to be passed if it is time-evolving.

Parameters

- grid** [Lattice object] Define the geometry of the simulation.
- state1** [State object] First component's state of the system.
- hamiltonian** [Hamiltonian object] Hamiltonian of the two-component system.
- delta_t** [float] A single evolution iteration, evolves the state for this time.
- Potential: Potential object, optional.** Time-evolving potential in component one.
- state2** [State object, optional.] Second component's state of the system.
- Potential2: Potential object, optional.** Time-evolving potential in component two.
- kernel_type** [str, optional (default: 'cpu')] Which kernel to use (either cpu or gpu).

Returns

- Solver** [Solver object] Solver object for the simulation of a two-component system.

Example

```

>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state_1 = ts.GaussianState(grid, 1.) # Create first-component system's
↪state
>>> state_2 = ts.GaussianState(grid, 1.) # Create second-component system's
↪state
>>> potential = ts.HarmonicPotential(grid, 1., 1.) # Create harmonic
↪potential
>>> hamiltonian = ts.Hamiltonian2Component(grid, potential, potential) #
↪Create an harmonic oscillator Hamiltonian
>>> solver = ts.Solver(grid, state_1, hamiltonian, 1e-2, State2=state_2) #
↪Create the solver

```

Members

evolve (*iterations*, *imag_time=False*)

Evolve the state of the system.

Parameters

- iterations** [integer] Number of iterations.
- imag_time** [bool,optional (default: False)] Whether to perform imaginary time evolution (True) or real time evolution (False).

Notes

The norm of the state is preserved both in real-time and in imaginary-time evolution.

Example

```

>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> potential = ts.HarmonicPotential(grid, 1., 1.) # Create harmonic
↪potential
>>> hamiltonian = ts.Hamiltonian(grid, potential) # Create a harmonic
↪oscillator Hamiltonian
>>> solver = ts.Solver(grid, state, hamiltonian, 1e-2) # Create the solver
>>> solver.evolve(1000) # perform 1000 iteration in real time evolution

```

get_inter_species_energy ()

Get the inter-particles interaction energy of the system.

Returns

- get_inter_species_energy** [float] Inter-particles interaction energy of the system.

get_intra_species_energy (*which=3*)

Get the intra-particles interaction energy of the system.

Parameters

- which** [integer,optional (default: 3)] Which intra-particles interaction energy to return: total system (default, which=3), first component (which=1), second component (which=2).

get_kinetic_energy (*which=3*)

Get the kinetic energy of the system.

Parameters

- which** [integer,optional (default: 3)] Which kinetic energy to return: total system (default, which=3), first component (which=1), second component (which=2).

get_potential_energy (*which=3*)
Get the potential energy of the system.

Parameters

- which** [integer,optional (default: 3)] Which potential energy to return: total system (default, which=3), first component (which=1), second component (which=2).

get_rabi_energy ()
Get the Rabi energy of the system.

Returns

- get_rabi_energy** [float] Rabi energy of the system.

get_rotational_energy (*which=3*)
Get the rotational energy of the system.

Parameters

- which** [integer,optional (default: 3)] Which rotational energy to return: total system (default, which=3), first component (which=1), second component (which=2).

get_squared_norm (*which=3*)
Get the squared norm of the state (default: total wave-function).

Parameters

- which** [integer,optional (default: 3)] Which squared state norm to return: total system (default, which=3), first component (which=1), second component (which=2).

get_total_energy ()
Get the total energy of the system.

Returns

- get_total_energy** [float] Total energy of the system.

Example

```
>>> import trottersuzuki as ts # import the module
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create the system's state
>>> potential = ts.HarmonicPotential(grid, 1., 1.) # Create harmonic_
↪potential
>>> hamiltonian = ts.Hamiltonian(grid, potential) # Create a harmonic_
↪oscillator Hamiltonian
>>> solver = ts.Solver(grid, state, hamiltonian, 1e-2) # Create the solver
>>> solver.get_total_energy() # Get the total energy
1
```

Solver::update_parameters()
Notify the solver if any parameter changed in the Hamiltonian

Tools

center_coordinates(*grid, x, y=None*):
Center and scale the coordinates of the grid to physical locations.

Parameters

- grid**: Lattice object Defines the topology.

- x: int.** Grid point.

- y: int, optional.** Grid point, 2D case.

Get the position of a single vortex in the quantum state.

get_vortex_position (*grid, state, approx_cloud_radius=0.*)

Get the position of a single vortex in the quantum state.

Parameters

- grid** [Lattice object] Define the geometry of the simulation.

- state** [State object] System's state.

- approx_cloud_radius** [float, optional] Radius of the circle, centered at the Lattice2D's origin, where the vortex core is expected to be. Need for a better accuracy.

Returns

- coords** [numpy array] Coordinates of the vortex core's position (coords[0]: x coordinate; coords[1]: y coordinate).

Notes

Only one vortex must be present in the state.

Example

```
>>> import trottersuzuki as ts # import the module
>>> import numpy as np
>>> grid = ts.Lattice2D() # Define the simulation's geometry
>>> state = ts.GaussianState(grid, 1.) # Create a state with gaussian wave_
↪function
>>> def vortex_a(x, y): # Define the vortex to be imprinted
>>>     z = x + 1j*y
>>>     angle = np.angle(z)
>>>     return np.exp(1j * angle)
>>> state.imprint(vortex) # Imprint the vortex on the state
>>> ts.get_vortex_position(grid, state)
array([ 8.88178420e-16,  8.88178420e-16])
```

D

delta_x (trottersuzuki.Lattice1D attribute), 14
 delta_x (trottersuzuki.Lattice2D attribute), 15
 delta_y (trottersuzuki.Lattice2D attribute), 15
 dim_x (trottersuzuki.Lattice1D attribute), 13
 dim_x (trottersuzuki.Lattice2D attribute), 15
 dim_y (trottersuzuki.Lattice2D attribute), 15

E

evolve() (trottersuzuki.Solver method), 32
 ExponentialState (class in trottersuzuki), 18
 ExponentialState() (trottersuzuki.ExponentialState method), 18

G

GaussianState (class in trottersuzuki), 21
 GaussianState() (GaussianState method), 21
 get_inter_species_energy() (trottersuzuki.Solver method), 32
 get_intra_species_energy() (trottersuzuki.Solver method), 32
 get_kinetic_energy() (trottersuzuki.Solver method), 32
 get_mean_px() (trottersuzuki.ExponentialState method), 19
 get_mean_px() (trottersuzuki.GaussianState method), 22
 get_mean_px() (trottersuzuki.SinusoidState method), 25
 get_mean_px() (trottersuzuki.State method), 16
 get_mean_pxp() (trottersuzuki.ExponentialState method), 19
 get_mean_pxp() (trottersuzuki.GaussianState method), 23
 get_mean_pxp() (trottersuzuki.SinusoidState method), 26
 get_mean_pxp() (trottersuzuki.State method), 17
 get_mean_py() (trottersuzuki.ExponentialState method), 20
 get_mean_py() (trottersuzuki.GaussianState method), 23
 get_mean_py() (trottersuzuki.SinusoidState method), 26
 get_mean_py() (trottersuzuki.State method), 17
 get_mean_pypy() (trottersuzuki.ExponentialState method), 20

get_mean_pypy() (trottersuzuki.GaussianState method), 23
 get_mean_pypy() (trottersuzuki.SinusoidState method), 26
 get_mean_pypy() (trottersuzuki.State method), 17
 get_mean_xx() (trottersuzuki.ExponentialState method), 20
 get_mean_xx() (trottersuzuki.GaussianState method), 23
 get_mean_xx() (trottersuzuki.SinusoidState method), 26
 get_mean_xx() (trottersuzuki.State method), 17
 get_mean_xx() (trottersuzuki.ExponentialState method), 20
 get_mean_xx() (trottersuzuki.GaussianState method), 23
 get_mean_xx() (trottersuzuki.SinusoidState method), 26
 get_mean_xx() (trottersuzuki.State method), 17
 get_mean_y() (trottersuzuki.ExponentialState method), 20
 get_mean_y() (trottersuzuki.GaussianState method), 23
 get_mean_y() (trottersuzuki.SinusoidState method), 26
 get_mean_y() (trottersuzuki.State method), 17
 get_mean_yy() (trottersuzuki.ExponentialState method), 20
 get_mean_yy() (trottersuzuki.GaussianState method), 23
 get_mean_yy() (trottersuzuki.SinusoidState method), 26
 get_mean_yy() (trottersuzuki.State method), 17
 get_particle_density() (trottersuzuki.ExponentialState method), 20
 get_particle_density() (trottersuzuki.GaussianState method), 23
 get_particle_density() (trottersuzuki.SinusoidState method), 26
 get_particle_density() (trottersuzuki.State method), 17
 get_phase() (trottersuzuki.ExponentialState method), 20
 get_phase() (trottersuzuki.GaussianState method), 23
 get_phase() (trottersuzuki.SinusoidState method), 26
 get_phase() (trottersuzuki.State method), 17
 get_potential_energy() (trottersuzuki.Solver method), 32
 get_rabi_energy() (trottersuzuki.Solver method), 33
 get_rotational_energy() (trottersuzuki.Solver method), 33
 get_squared_norm() (trottersuzuki.ExponentialState method), 20
 get_squared_norm() (trottersuzuki.GaussianState method), 23

method), 23
get_squared_norm() (trottersuzuki.SinusoidState method), 26
get_squared_norm() (trottersuzuki.Solver method), 33
get_squared_norm() (trottersuzuki.State method), 17
get_total_energy() (trottersuzuki.Solver method), 33
get_value() (trottersuzuki.HarmonicPotential method), 29
get_value() (trottersuzuki.Potential method), 28
get_vortex_position(), 34
get_x_axis() (trottersuzuki.Lattice1D method), 13
get_x_axis() (trottersuzuki.Lattice2D method), 14
get_y_axis() (trottersuzuki.Lattice2D method), 14

H

Hamiltonian (class in trottersuzuki), 29
Hamiltonian() (Hamiltonian method), 29
Hamiltonian2Component (class in trottersuzuki), 30
Hamiltonian2Component() (Hamiltonian2Component method), 30
HarmonicPotential (class in trottersuzuki), 28

I

imprint() (trottersuzuki.ExponentialState method), 19
imprint() (trottersuzuki.GaussianState method), 22
imprint() (trottersuzuki.SinusoidState method), 25
imprint() (trottersuzuki.State method), 16
init_potential() (trottersuzuki.Potential method), 28

L

Lattice1D (class in trottersuzuki), 13
Lattice2D (class in trottersuzuki), 14
Lattice2D() (Lattice1D method), 13
Lattice2D() (Lattice2D method), 14
length_x (trottersuzuki.Lattice1D attribute), 13
length_x (trottersuzuki.Lattice2D attribute), 15
length_y (trottersuzuki.Lattice2D attribute), 15
loadtxt() (trottersuzuki.ExponentialState method), 21
loadtxt() (trottersuzuki.GaussianState method), 24
loadtxt() (trottersuzuki.SinusoidState method), 27
loadtxt() (trottersuzuki.State method), 18

P

Potential (class in trottersuzuki), 27
Potential() (Potential method), 27

S

SinusoidState (class in trottersuzuki), 24
SinusoidState() (SinusoidState method), 24
Solver (class in trottersuzuki), 31
State (class in trottersuzuki), 15
State() (State method), 15

W

write_particle_density() (trottersuzuki.ExponentialState method), 21
write_particle_density() (trottersuzuki.GaussianState method), 24
write_particle_density() (trottersuzuki.SinusoidState method), 27
write_particle_density() (trottersuzuki.State method), 18
write_phase() (trottersuzuki.ExponentialState method), 21
write_phase() (trottersuzuki.GaussianState method), 24
write_phase() (trottersuzuki.SinusoidState method), 27
write_phase() (trottersuzuki.State method), 18
write_to_file() (trottersuzuki.ExponentialState method), 21
write_to_file() (trottersuzuki.GaussianState method), 24
write_to_file() (trottersuzuki.SinusoidState method), 27
write_to_file() (trottersuzuki.State method), 18